主要参考文献

[1] 郭佩霞、徐家聪，1980：对安徽巢县青龙群时代的认识，地层学杂志，4卷，4期。

[2] 全国地层委员会，1981：中国地层指南及中国地层指南说明书。科学出版社。

[3] 王志浩、曹延岳，1981：湖北利川早三叠世牙形刺。古生物学报，20卷，4期。

[4] 阿格尔，D.V.（王仪诚译），1981：古生态学原理。科学出版社。

[5] 奥德姆，E.P.（孔儒泳等译），1981：生态学基础。人民教育出版社。

[6] 闵庆魁、陈楚震，1981：南京地区三叠系的再研究。地层学杂志，5卷，1期。

[7] 张仁杰，1982：湖北东南部三叠纪地层。宜昌地质矿产研究所所刊，5号。

[8] 郭佩霞，1982：苏、皖早三叠世晚期菊石的发现。古生物学报，21卷，5期。

[9] 丁梅华，1983：安徽巢县马家山下三叠统的研究。地球科学，2期

[10] 杨式傅，1984：Zoophycos在中国的发现及其环境意义。石油天然气地质，5卷3期。

[11] 张正坤，1984：中朝地块与扬子地块在古生代晚期是太平洋古陆的一部分。中国地质科学院院报，第九号。

[12] 杨式傅，1985：我国遗迹化石研究新进展。地质论评，31卷3期。

[13] 林金录，1985：华南与华北断块的地极移动曲线。中国地质科学院院报、第九号。

[14] 殷鸿福、凌秋贤，1986：中国三叠纪生物地理区系。中国古生物学会第十三、十四届学术年会论文选集，中国古生物学会编辑，安徽科学技术出版社。

[15] 刘宝君，1986：四川兴文四龙下二叠统碳酸盐风景岩。地质学报60卷1期。

[16] 张国栋，1987：下扬子区早三叠世碳酸盐风暴流与碎屑流沉积特征。海洋地质与第四纪地质，7卷2期。

[17] 刘家润，张永辂，1987：滇东盘溪中泥盆世一个群落的分析一兼论样方法中化石计数问题。古生物学报，26卷，1期。

[18] 毕德昌，1987：试论地层单位的划分。中国区域地质，第1期。

[19] 冯增昭主编，1988：下扬子地区中下三叠统青龙群岩相古地理研究。云南科技出版社。

[20] 王义刚，1988：中国三叠纪菊石的古生态学概论。古生物学报，28卷，第3期。

[21] 张守信，1989：理论地层学——现代地层学概念。科学出版社。

[22] 王润华、薛虎，1989：下扬子地区中生界古地磁及其构造解析。南京地质矿产研究所所刊，第十卷，第一号。

[23] 吴汉宁，常承法等，1990：依据古地磁资料探讨华北和华南块体运动及其对秦岭造山带构造演化的影响。地质科学，第3期。

[24] Purdy, E.G., 1963: Recent calcium Carbonate facies of the Great Bahamas Bank, J.Geol., 71,334–472.

[25] Fagerstrom, J.A., 1964: Fossil communites in Paleoecology: Their recognition and significance. Geol. Soc. America Bull. Vol. 75, No.12.

[26] Tozer, E.T., 1966: A standard for Triasic Time. Geological survey of Canada Bulletin 156.

[27] Ginsburg, R.N.Hardie, L.A., 1975: Tidal and storm deposits, northeastern Andros Islands, Bahamas. In: Tidal Deposits: Acasebook of recent examples and fossil counterparts (Ed. by R.N., Ginsburg) 201–208.

[28] Frey, R.W., 1975: The study of trace fossils. Springer–Verlag, New York.

[29] Hantzschel, W., 1975: Trace Fossils and Problematica: Treatise on invertebrate Palaeontology, Pt. W., Geol.

Soc. Amer. and Univ. Kansas Press.

[30] Wilson, J.L., 1975: Carbonate Facies in Geologic History. Springer–Verlag, New York.

[31] Hollis, D.Hedberg, (ed.), 1976: A Guide to Stratigraphic Classification Teminology and Procedure. John Wiley and Sons Inc. New York.

[32] Dodd, J.R. and Stanton, R.J.Jr., 1981: Paleoecology, Concepts and Applications. John Wiley and Sons Inc. New York.

[33] Lehman, V., 1981: The Ammonites: Their Life and World. Combridge Univ. Press., London.

[34] Aigner, T., 1982: Calcareous Tempestites: Storm–dominated Stratification in Upper Muschelkalk Limestones (Middle Trias, SW–Germeny). in Cyclic and Event Stratification (ed. by Einsele / Seilacher).

[35] Dott, R.H.Jr., 1982: Hummocky Stratification Significance of its Variable Bedding Sequences. Geal. Soc. of Amer. Bull. Vol.93, No.8.

[36] Jeffery, D., and Aigner, T., 1982: Storm Sedimentation in the Carboniferous Limestones Near Weston–Super–Mare (Dinantian, SW–England). in Cyclic and Event Stratification (ed. by Einsele / Seilacher).

[37] Ekdale, A.A., Bromley, R.G., and Pemberton, S.G., 1984: Ichnology: The use of Trace Fossils in Sedimentology and Stratigraphy. Soc. of Econ. Paleont. and Mineral. Tulss. Oklahoma.

[38] Curran, H.A., 1985: Biogenic Structures: Their use in interpreting depositional environments. Soc. Econ. Paleont. Mineral. Special Publication. No.35.

[39] Aigner, T., 1985: Storm depositional Systems. in Lecture Notes in Earth Sciences 3.Springer–Verlag.

[40] Read, J.F., 1985: Carbonate Platform Facies Models. A.A.P.G.Vol. 69 No.1.